

## Distribution of the bloom-forming diatom *Didymosphenia geminata* in the Ebro River basin (Northeast Spain) in the period 2006-2009

Pedro Tomás<sup>1\*</sup>, Javier Oscoz<sup>2</sup>, Concha Durán<sup>3</sup>, David Fernández<sup>4</sup> and José Pedro Marín<sup>5</sup>

<sup>1</sup>Lab. de Ensayos Técnicos S.A., Polig. Ind. Valdeconsejo, C/ Aneto, Parcela 8-A, E-50410 Cuarte de Huerva, Zaragoza, Spain

<sup>2</sup>Dep. of Zoology and Ecology, Faculty of Sciences, University of Navarra, P.O. Box 177, E-31080, Pamplona, Navarra, Spain

<sup>3</sup>Área Calidad de Aguas, Confederación Hidrográfica del Ebro, Paseo Sagasta 24-28, E-50071, Zaragoza, Spain

<sup>4</sup>Hydraena s.l.l., Plaza del Baño n°4, E-18170, Alfacar, Granada, Spain

<sup>5</sup>Dpto. de Didáctica de las Ciencias Experimentales, Univ. de Murcia, Campus de Espinardo, E-30100, Espinardo, Murcia, Spain

E-mail: [biologia@ensaya.es](mailto:biologia@ensaya.es) (PT), [joscoz@alumni.unav.es](mailto:joscoz@alumni.unav.es) (JO), [cduran@chebro.es](mailto:cduran@chebro.es) (CD), [hydraena@hydraena.com](mailto:hydraena@hydraena.com) (DF),

[jpmurcia@um.es](mailto:jpmurcia@um.es) (JPM)

\*Corresponding author

Received: 17 December 2009 / Accepted: 2 August 2010 / Published online: 23 August 2010

### Abstract

This paper presents the distribution and the ecological parameters of *Didymosphenia geminata* in the Ebro River basin (Northeast Spain) during the period 2006-2009. The presence of blooms in some rivers in 2009, as in Cinca River in Las Pilas bridge, is reported. *D. geminata* was found in summer at 24 localities, forming blooms in three of them. It prefers mountainous calcareous rivers, regulated or not, with oligotrophic waters, low flow conditions and well illuminated stretches. In order to prevent nuisance blooms of the species, it is necessary to study its ecology and establish a monitoring network for detecting early stages of blooms in order to take control measures.

**Key words:** *Didymosphenia geminata*, diatom, ecological parameters, nuisance organism, Ebro basin

### Introduction

The algae, *Didymosphenia geminata* (Lyngbye) Schmidt, 1899, is a freshwater stalked diatom characterized by having large triundulate frustules (Figure 1). *D. geminata* (didymo) forms fibrous masses of yellow-brown, more than 3 cm thick, like wool or cellulose aggregates, although its colour varies depending on the type of sediments retained (Kilroy 2004). It was originally described in the Faroe Islands in 1819 by Lyngbye, and its fossil record and biogeographic studies show that its original distribution only covered the Northern Hemisphere, always at latitudes above 30°N (Blanco and Ector 2009). However, different studies showed that didymo is expanding its ecological range in environmental conditions (Kawecka and Sanecki 2003; Kilroy 2004; Spaulding and Elwell 2007). In the last few decades it has occurred more and more frequently across worldwide freshwater ecosystems and it has been cited in over 50 countries.


Didymo blooms were reported in their areas of origin in the 19th century (e.g. in the British Isles) and it seemed to be somewhat common (Whitton et al. 2009). However, blooms are increasing in Eurasian and North American areas, where they had not formed before, and these have caught the attention of researchers in the last decades. There are many studies and theories about factors leading to the formation of massive growth, for instance regulation of rivers, genetic variation, climate change, changes in land use in the watershed or changes in the N: P balance (see Whitton et al. 2009). These blooms can cover the substrate of the riverbed along several kilometres, so that this diatom is likely to have significant impacts on native species and biodiversity through ecological processes (Campbell 2005; Shearer and Erickson 2006; Larned et al. 2007; Spaulding and Elwell 2007; Larson 2008; Gillis and Chalifour 2010). Didymo is considered invasive in several countries, as in New Zealand where it was declared as an invasive nuisance in 2004.

In the Iberian Peninsula this species has been found in some localities of the Duero River basin (Blanco and Bécares 2009) and there are historical records of its presence in the Pyrenees and Mallorca (Margalef 1953; URS 2006). A complete list of records can be found in Blanco and Ector (2009). The first observations of massive growth of *D. geminata* in the Ebro River basin was in 2005 in the Ara River upstream of Ainsa, but it was not confirmed that they were due to *D. geminata* until the summer of 2006, with the new appearance of the species mass, (URS 2006). In 2006, it was also recorded from the Subordán River in Hecho (Ortiz and Cambra 2006). In subsequent years it has appeared in low abundance in several samples of biological control networks of the Confederación Hidrográfica del Ebro, although it is noted that the sampling protocol applied, underestimates the presence of this diatom (Kirkwood et al. 2007).

The objective of this study is to contribute to the knowledge of the presence and formation of blooms of *D. geminata* in the Ebro River basin, providing new data on its distribution in this area.


### Materials and methods

A description of the Ebro River basin can be found in Oscoz et al. (2010). During summer biological sampling of the CEMAS network to assess the water quality of the Ebro River Basin in the period 2006-2009, 2006 (n= 211), 2007 (n= 323), 2008 (n= 343), 2009 (n= 362), a total of 1239 stations, some of them were visited annually. In each station samples of macro-invertebrates, diatoms and macrophytes were taken. The sampling and the analysis of the diatom samples were done according to CHE protocols (2005). At least five cobbles or small boulders were collected from unshaded areas within the main flow and free from obvious filamentous algae or siltation, and the upper surface of the stones was brushed with a toothbrush to remove the diatom film. The toothbrush was rinsed in a container with water. The sample was fixed with 4% formalin. In the literature (Spaulding and Elwell 2007) it is stated that this sampling and analytical method is inadequate to pick it up, therefore the valves detected in the samples of macrophytes were also recorded. In the laboratory the samples were oxidized with hydrogen peroxide and mounted on a slide with naphrax, for more details see the protocol. At least 400 valves were counted.



**Figure 1.** *Didymosphenia geminata* from Gállego River (code 0808 in Sta. Eulália) (Scale Bar: 10µm, photograph by JPM).

Physico-chemical parameters (pH, temperature, conductivity and dissolved oxygen) were obtained in situ with a multiparametric sonde YSI 556MPS. Water samples were taken and, according to the methodologies established in the ITC-MMA.EECC-1/06 (Ministry of the Environment 2006), they were conserved at 4°C in order to analyse the following parameters in the laboratory: nitrates, nitrites, phosphates and silica. Other hydromorphological features such as regulation by dams, flow and turbulence, type of substrata and illumination of the riverbed were assessed qualitatively, according to the ranges established in Pardo et al. (2002). After sampling all the used material was disinfected according to the protocols of disinfection of the Ebro River basin to prevent the spread of the zebra mussel (*Dreissena polymorpha* (Pallas, 1771)) and other organisms (CHE 2007). All sampling was carried out following a stated period of steady flows (four weeks).



**Figure 2.** Distribution of *Didymosphenia geminata* in the Ebro River basin, 2006-2009. black points: presence, grey points: absence, white points: bloom.

## Results and discussion

*Didymosphenia geminata* was recorded in 24 sampling stations in the studied years (Figure 2, Annex 1). The number of localities where didymo was presented varied in the different years: Two stations in 2006, nine stations in 2007, eight localities in 2008 and 16 sample stations in 2009. It was recorded preferentially in calcareous mountainous rivers, and only in three localities did this species form blooms (Cinca River in Puente de las Pilas, Cinca River in Ainsa and Subordan River in Embún). The distribution of the diatom in the Ebro River basin is presented in the Figure 2. Most of the localities where didymo was recorded were located in Pyrenean rivers. Only in the locality of Almarza, Iregua River (code 1184) was not located in this area. The presence of didymo in the headwaters of the Iregua River could be due to the accidental introduction from the nearby Vinuesa River (Duero River Basin) where the species is present forming blooms of varying length, both as clumps of didymo as well as

single microscopic cells, which can be moved around by boaters, anglers, hunters and other aquatic recreationists.

In eight localities didymo was present for at least two years; and it can be noted that at the Ara River station in Ainsa it was found in all years. In general, the abundance of the species in the samples was low (Annex 1). This low abundance could seem contradictory in the case of the localities where blooms were recorded, but it was due to the fact that the diatom samples were taken in boulders and cobbles free of didymo blooms.

According to our data on ecological parameters didymo in Ebro River basin was found in well oxygenated waters (8.2-12.0 mg/l dissolved O<sub>2</sub>), with a pH range of 7.3 to 9.0, and water temperature ranging from 11.6-24.7°C. Conductivity was relatively medium-low (117-511 µs/cm) and the concentration of nutrients was low, NO<sub>3</sub> (<0.63-3.67 mg/l), NO<sub>2</sub> (0.01-0.04 mg/l), NH<sub>4</sub> (<0.01-0.28 mg/l), PO<sub>4</sub> (<0.05-0.73 mg/l); probably because didymo was recorded in mountain streams, mainly in upper oligotrophic

river stretches where these parameters are usually lower. These rivers were located in an altitude range between 337 and 1186 meters.

According to habitat characteristics this species was mainly found under low-medium flow conditions and low-high water turbulence, both in high and low turbidity waters, although it was found mainly in well illuminated river stretches. The substrate of the rivers where didymo was present was composed mainly of cobbles and boulders. Some authors suggested a relationship between the presence of the diatom blooms and regulated or lake fed streams (Kilroy et al. 2006; Kawecka and Sanecki, 2003; Kirkwood et al. 2007), but according to our data the regulation of the river could not affect the presence of *D. geminata*, as it has been recorded forming blooms both in regulated rivers (as Cinca River) and in unregulated rivers (such as the Ara River). All of the measured ecological and physico-chemical parameters measured in our study were in accordance with the ranges found in the literature according to Campbell (2005) and Whitton et al. (2009).

## Conclusions

It seems that *Didymosphenia geminata* is expanding its distribution in the Ebro River basin in the last few years or, according to the literature (Whitton et al. 2009), some years the habitat conditions or other unknown parameters favoured the growth of this diatom, however the presence of blooms is not usually frequent. According to our data this species prefers mountainous calcareous rivers, regulated or not, with oligotrophic waters, relatively low flow conditions and well illuminated stretches. From our data it is difficult to relate the presence or absence of didymo only to physical and chemical characteristics of the water, as it appears to be present in a wide range of environmental conditions, this means that the potential for spread is great because the environment is not a limiting factor. The presence of *D. geminata* in the Iregua River is probably due to traslocation from the Vinuesa River may indicate the need for increasing public information and awareness about disinfection protocols (as in zebra mussel (*Dreissena polymorpha*) it is important to check, clean and dry). In order to prevent the nuisance blooms of the species, it is also necessary to study the ecology and to establish a monitoring network for detect early stages of the blooms in

order to take measures of control when these early stages are detected.

## Acknowledgements

Authors are grateful to all people that collaborate in the different sampling carried out in the Ebro River basin in the last number of years. Thanks to R. Ortiz, N. Flor and J. Cambra for the data of their surveys. Thanks also to Andrés Mellado, Isidoro Pérez, José Miguel García, Pablo Jáimez-Cuellar, Julio M. Luzón, José Antonio Palomino, for their work in the Ebro River basin surveys during 2006-2009.

## References

- Blanco S, Ector L (2009) Distribution, ecology and nuisance effects of the freshwater invasive diatom *Didymosphenia geminata* (Lyngbye) M. Schmidt: a literature review. *Nova Hedwigia* 88: 347–422, doi:10.1127/0029-5035/2009/0088-0347
- Blanco S, Bécarea E (2009) Una nueva alga invasora llega a España: *Didymosphenia geminata*. *Quercus* 283:30–35
- Campbell ML (2005) Organism Impact Assessment (OIA) for potential Impacts of *Didymosphenia geminata*. All Oceans Ecology. Australia, 92 pp
- CHE, Confederación Hidrográfica del Ebro (2005) Metodología para el establecimiento del Estado Ecológico según la Directiva Marco del Agua. Protocolos de muestreo y análisis. Ministerio de Medio Ambiente, 43 pp
- CHE, Confederación Hidrográfica del Ebro (2007) Protocolo de desinfección de embarcaciones y equipos en masas de agua infectadas por mejillón cebra (*Dreissena polymorpha*). Ministerio de Medio Ambiente y Medio Rural y Marino, 12 pp
- Gillis CA, Chalifour M (2010) Changes in the macrobenthic community structure following the introduction of the invasive algae *Didymosphenia geminata* in the Matapedia River (Québec, Canada). *Hydrobiologia* 647: 63–70, doi:10.1007/s10750-009-9832-7
- ITC-MMA.EECC-1/06 (2006) ORDEN MAM/3207/2006, de 25 de septiembre, por la que se aprueba la instrucción técnica complementaria MMA-EECC-1/06, determinaciones químicas y microbiológicas para el análisis de las aguas. BOE 250: 36326–36440
- Kawecka B, Sanecki J (2003) *Didymosphenia geminata* in running waters of southern Poland – symptoms of change in water quality? *Hydrobiologia* 495: 193–201, doi:10.1023/A:1025469500265
- Kilroy C (2004) A new alien diatom, *Didymosphenia geminata* (Lyngbye) Schmidt – its biology, distribution, effects and potential risks for New Zealand fresh waters. National Institute of Water and Atmospheric Research. New Zealand. Client Report, CHC2004-128, 40 pp
- Kilroy C, Lagerstedt A, Davey A, Robinson K (2006) Studies on the survivability of the exotic, invasive diatom *Didymosphenia geminata* under a range of environmental and chemical conditions. National Institute of Water and Atmospheric Research. New Zealand. Client Report CHC2006-116, NIWA Project MAF06506, 121 pp
- Kirkwood A, Shea ET, Jackson LJ, McCauley E (2007) *Didymosphenia geminata* in two Alberta headwater rivers: an emerging invasive species that challenges conventional views on algal bloom development. *Canadian Journal of Fisheries and Aquatic Sciences* 64: 1703–1709, doi:10.1139/F07-152

Distribution of *Didymosphenia geminata* in the Ebro River basin

- Larned S, Biggs B, Blair N, Burns C, Jarvie B, Jellyman D, Kilroy C, Leathwick J, Lister K, Nagels J, Schallenberg M, Sutherland S, Sykes J, Thompson W, Vopel K, Wilcock B (2007) Ecological studies of *Didymosphenia geminata* in New Zealand, 2006-2007. National Institute of Water and Atmospheric Research. New Zealand. Client Report CHC2007-070, NIWA Project MAF07507, 140 pp
- Larson AM, Carreiro J (2008) Relationship between nuisance blooms of *Didymosphenia geminata* and measures of aquatic community composition in Rapid Creek, South Dakota. In: Bothwell ML, Spaulding SA (eds) Proceedings of the 2007 International Workshop on *Didymosphenia geminata*. *Canadian Technical Report on Fisheries and Aquatic Sciences* 2795: 45-49
- Margalef R (1953) Materiales para la hidrobiología de la Isla de Mallorca. *Publicaciones del Instituto Biología Aplicada Barcelona* 15: 5-111
- Ortiz R, Cambra J (2006) Red de Intercalibración, Red de Referencia y Red Básica de diatomeas en la Cuenca del Ebro, Informe Final 2005-2006. Universitat de Barcelona, 93 pp
- Oscosz J, Tomás P, Durán C (2010). Review and new records of non-indigenous freshwater invertebrates in the Ebro River basin (Northeast Spain). *Aquatic Invasions* 5: 263-284, doi: 10.3391/ai.2010.5.3.04
- Pardo I, Álvarez M, Casas J, Moreno JL, Vivas S, Bonada N, Alba-Tercedor J, Jáimez- Cuéllar P, Moyà G, Prat N, Robles S, Suárez ML, Toro M, Vidal-Albarca MR (2002) El hábitat de los ríos mediterráneos. Diseño de un índice de diversidad de hábitat. *Limnetica* 21(3-4): 115-133
- Shearer J, Erickson J (2006) *Didymosphenia geminata* and the Rapid Creek brown trout fishery, South Dakota. In: Spaulding S, Wiltshire R, Elwell L (eds), Current Knowledge of *Didymosphenia geminata*: Developing a Research and Management Response. Federation of Fly Fishers and EPA Region 8, held in association with Western Division American Fisheries Society Annual Meeting, May 15-16, 2006. Montana State University Montana. USA. pp 22
- Spaulding S, Elwell L (2007) Increase in nuisance blooms and geographic expansion of the freshwater diatom *Didymosphenia geminata*. Environmental Protection Agency and the Federation of Fly Fishers Montana: 33 pp
- URS (2006) Investigación del origen de depósitos sobre sustratos sumergidos, en el río Ara (cuenca del Cinca). Confederación Hidrográfica del Ebro. Barcelona, 30 pp
- Whitton BA, Ellwood NTW, Kawecka B (2009) Biology of the freshwater diatom *Didymosphenia*: a review. *Hydrobiologia* 630:1-37, doi:10.1007/s10750-009-9753-5

**Annex 1.** Records of *Didymosphenia geminata* in the Ebro River basin in the period 2006-2009. Abundance in n° of valves in the diatom samples (\* found in the sample outside the count area).

Code	River/Locality	Geographic coordinates		Abundance			
		Latitude, N	Longitude, W	2006	2007	2008	2009
0013	Ésera / Graus	42°12'24"	0°20'49"	-	3	-	-
0018	Aragón / Jaca	42°34'45"	0°33'10"	-	*	-	-
0101	Aragón / Yesa	42°37'02"	1°12'34"	-	-	*	*
0561	Gállego / Jabarrella	42°24'12"	0°29'51"	-	-	-	*
0616	Cinca / Derivación Acequia Paules	42°02'29"	0°11'10"	-	*	-	-
0802	Cinca / Puente de las Pilas, Estada-Estadilla	42°03'53"	0°12'58"	-	-	*	*
0808	Gállego/Santa Eulàlia	42°16'12"	0°45'03"	-	-	-	*
0804	Subordán / La Peñeta	42°44'17"	0°44'46"	*	4	-	-
0816	Esca/Burgui	42°43'04"	1°00'15"	-	-	*	-
1056	Veral / Biniés	42°37'37"	0°49'08"	-	12	-	-
1088	Gállego/Biescas	42°37'41"	0°19'15"	-	-	*	*
1089	Gállego / Sabinánigo	42°30'41"	0°21'20"	-	-	*	-
1090	Gállego / Hostal de Ipiés	42°26'35"	0°23'39"	-	*	-	*
1092	Gállego / Murillo de Gállego	42°19'13"	0°45'14"	-	-	-	*
1121	Cinca / Laspuña	42°30'25"	0°08'54"	-	*	1	*
1122	Cinca / Aínsa	42°24'57"	0°08'39"	-	-	-	*
1130	Ara / Torla E.A. 196	42°37'52"	0°06'25"	-	2	-	*
1131	Ara / Fiscal	42°30'00"	0°06'56"	-	-	-	*
1132	Ara / Aínsa	42°24'52"	0°08'29"	*	3	1	*
1184	Iregua/Almarza	42°13'39"	2°37'33"	-	-	-	*
2010	Irati/Lumbier	42°40'03"	1°19'00"	-	-	*	-
2024	Subordán / Embún	42°37'37"	0°43'14"	-	-	-	*
2199	Escarra/Escarrilla	42°44'17"	0°19'03"	-	-	-	*
2211	Bellós / Puyarruego	42°33'44"	0°03'05"	-	-	-	*